

References:

[Bilbao-Calabuig R et al. Visual outcomes following bilateral implantation of two diffractive trifocal intraocular lenses in 10,084 eyes. American Journal of Ophthalmology, July 2017.](#)

[Ang R. Long Term Clinical Outcomes of Hydrophilic and Hydrophobic Versions of a Trifocal IOL with the same optical design. Clinical Ophthalmology. 2023;17.](#)

[Poyales F, Pérez R, López-Brea I, Zhou Y, Rico L, Garzón N. Comparison of Visual Performance and Patient Satisfaction Outcomes with Two Trifocal IOLs with Similar Optical Design but Different Materials. Clin Ophthalmol. 2020;14:3237-3247.](#)

[Chassain C, Chamard C. Posterior capsule opacification, glistenings and visual outcomes: 3 years after implantation of a new hydrophobic. Journal Français d'Ophtalmologie 2018; 513-520.](#)

[Ang RET, Tañá-Rivero P, Pastor-Pascual F, Stodulka P, Tetz M, Fischinger I. Visual and Refractive Outcomes After Bilateral Implantation of a Biconvex Aspheric Toric Monofocal Intraocular with a Double C-Loop Haptic Design. Clinical Ophthalmology 2023;17 2765–2776.](#)

[Bozukova D, Pagnouille C, Jérôme C. Biomechanical and optical properties of 2 new hydrophobic platforms for intraocular lenses. J Cataract Refract Surg 2013 Sep;39\(9\):1404-14.](#)